

**OUTPUT 4: APPENDIX-II SPECIES WITH MINIMAL (<= 5 TRADE RECORDS)
OR NO TRADE OVER THE PERIOD 1999-2009
(based on number of trade records and excluding trade in sources C and D)**

| Order | Family | Taxon | No. trade records | | IUCN |
|------------------------|--------------|------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| PHYLUM CHORDATA | | | | | |
| CLASS MAMMALIA | | | | | |
| ARTIODACTYLA | BOVIDAE | <i>Cephalophus brookei</i> | 0 | 0 | NT |
| | | <i>Saiga borealis</i> | 0 | 0 | |
| | CERVIDAE | <i>Cervus elaphus bactrianus</i> | 3 | 1 | LC |
| | | <i>Pudu mephistophiles</i> | 1 | 1 | VU |
| CARNIVORA | CANIDAE | <i>Cuon alpinus</i> | 1 | 0 | EN |
| | | <i>Lycalopex fulvipes</i> | 1 | 1 | CR |
| | | <i>Vulpes cana</i> | 5 | 2 | LC |
| | MUSTELIDAE | <i>Lutra sumatrana</i> | 0 | 0 | EN |
| | OTARIIDAE | <i>Arctocephalus galapagoensis</i> | 1 | 1 | EN |
| | | <i>Arctocephalus philippii</i> | 1 | 1 | NT |
| | VIVERRIDAE | <i>Cynogale bennettii</i> | 4 | 4 | EN |
| | | <i>Prionodon linsang</i> | 2 | 2 | LC |
| CHIROPTERA | PTEROPODIDAE | <i>Acerodon celebensis</i> | 2 | 1 | LC |
| | | <i>Acerodon humilis</i> | 3 | 3 | EN |
| | | <i>Acerodon leucotis</i> | 0 | 0 | VU |
| | | <i>Acerodon mackloti</i> | 0 | 0 | VU |
| | | <i>Pteropus admiralitatum</i> | 2 | 2 | LC |
| | | <i>Pteropus aldabrensis</i> | 1 | 0 | VU |
| | | <i>Pteropus alecto</i> | 3 | 3 | LC |
| | | <i>Pteropus anetianus</i> | 3 | 0 | VU |
| | | <i>Pteropus aruensis</i> | 0 | 0 | CR |
| | | <i>Pteropus banakrisi</i> | 0 | 0 | |
| | | <i>Pteropus brunneus</i> | 0 | 0 | EX |
| | | <i>Pteropus caniceps</i> | 0 | 0 | NT |
| | | <i>Pteropus capistratus</i> | 0 | 0 | NT |
| | | <i>Pteropus chrysoproctus</i> | 0 | 0 | NT |
| | | <i>Pteropus cognatus</i> | 0 | 0 | EN |
| | | <i>Pteropus conspicillatus</i> | 3 | 3 | LC |
| | | <i>Pteropus dasymallus</i> | 4 | 3 | NT |
| | | <i>Pteropus faunulus</i> | 0 | 0 | VU |
| | | <i>Pteropus fundatus</i> | 0 | 0 | EN |
| | | <i>Pteropus gilliardorum</i> | 0 | 0 | DD |
| | | <i>Pteropus griseus</i> | 0 | 0 | DD |
| | | <i>Pteropus howensis</i> | 0 | 0 | DD |
| | | <i>Pteropus intermedius</i> | 0 | 0 | DD |
| | | <i>Pteropus keyensis</i> | 0 | 0 | DD |
| | | <i>Pteropus leucopterus</i> | 1 | 1 | LC |
| | | <i>Pteropus lombocensis</i> | 0 | 0 | DD |
| | | <i>Pteropus lylei</i> | 2 | 2 | VU |
| | | <i>Pteropus mahaganus</i> | 0 | 0 | VU |

| Order | Family | Taxon | No. trade records | | IUCN |
|----------------|----------------|-----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Pteropus melanopogon</i> | 0 | 0 | EN |
| | | <i>Pteropus melanotus</i> | 0 | 0 | VU |
| | | <i>Pteropus neohibernicus</i> | 2 | 2 | LC |
| | | <i>Pteropus nitendiensis</i> | 0 | 0 | EN |
| | | <i>Pteropus ocularis</i> | 0 | 0 | VU |
| | | <i>Pteropus ornatus</i> | 1 | 1 | VU |
| | | <i>Pteropus personatus</i> | 0 | 0 | LC |
| | | <i>Pteropus pohlei</i> | 0 | 0 | EN |
| | | <i>Pteropus poliocephalus</i> | 3 | 3 | VU |
| | | <i>Pteropus pselaphon</i> | 0 | 0 | CR |
| | | <i>Pteropus rayneri</i> | 1 | 1 | NT |
| | | <i>Pteropus rennelli</i> | 0 | 0 | VU |
| | | <i>Pteropus scapulatus</i> | 1 | 1 | LC |
| | | <i>Pteropus speciosus</i> | 0 | 0 | DD |
| | | <i>Pteropus subniger</i> | 0 | 0 | EX |
| | | <i>Pteropus temminckii</i> | 0 | 0 | VU |
| | | <i>Pteropus tokudae</i> | 0 | 0 | EX |
| | | <i>Pteropus tuberculatus</i> | 0 | 0 | CR |
| | | <i>Pteropus vetulus</i> | 2 | 2 | VU |
| | | <i>Pteropus voeltzkowi</i> | 1 | 1 | VU |
| | | <i>Pteropus woodfordi</i> | 0 | 0 | VU |
| DIPROTODONTIA | MACROPODIDAE | <i>Dendrolagus inustus</i> | 3 | 3 | VU |
| | | <i>Dendrolagus ursinus</i> | 2 | 0 | VU |
| | PHALANGERIDAE | <i>Phalanger intercastellanus</i> | 0 | 0 | LC |
| | | <i>Phalanger mimicus</i> | 2 | 2 | LC |
| | | <i>Spilocuscus kraemeri</i> | 0 | 0 | NT |
| | | <i>Spilocuscus papuensis</i> | 0 | 0 | VU |
| MONOTREMATA | TACHYGLOSSIDAE | <i>Zaglossus attenboroughi</i> | 1 | 1 | CR |
| | | <i>Zaglossus bartoni</i> | 1 | 1 | CR |
| | | <i>Zaglossus bruijni</i> | 3 | 3 | CR |
| PERISSODACTYLA | EQUIDAE | <i>Equus kiang</i> | 3 | 0 | LC |
| PHOLIDOTA | MANIDAE | <i>Manis crassicaudata</i> | 2 | 0 | NT |
| | | <i>Manis culionensis</i> | 1 | 1 | NT |
| | | <i>Manis gigantea</i> | 3 | 3 | NT |
| PRIMATES | AOTIDAE | <i>Aotus azarae infulatus</i> | 0 | 0 | LC |
| | | <i>Aotus hershkovitzi</i> | 0 | 0 | VU |
| | | <i>Aotus lemurinus brumbacki</i> | 0 | 0 | VU |
| | | <i>Aotus miconax</i> | 0 | 0 | VU |
| | | <i>Aotus nigriceps</i> | 1 | 0 | LC |
| | ATELIDAE | <i>Alouatta belzebul</i> | 1 | 1 | VU |
| | | <i>Alouatta guariba</i> | 2 | 0 | LC |
| | | <i>Alouatta macconnelli</i> | 0 | 0 | LC |
| | | <i>Alouatta nigerrima</i> | 0 | 0 | LC |
| | | <i>Alouatta sara</i> | 0 | 0 | LC |
| | | <i>Ateles hybridus</i> | 0 | 0 | CR |
| | | <i>Ateles marginatus</i> | 1 | 0 | EN |
| | | <i>Lagothrix cana</i> | 0 | 0 | EN |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|-----------------|------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Lagothrix lugens</i> | 0 | 0 | CR |
| | CEBIDAE | <i>Callithrix acariensis</i> | 0 | 0 | |
| | | <i>Callithrix argentata</i> | 3 | 1 | |
| | | <i>Callithrix chrysoleuca</i> | 0 | 0 | DD |
| | | <i>Callithrix emiliae</i> | 0 | 0 | |
| | | <i>Callithrix humeralifera</i> | 3 | 0 | |
| | | <i>Callithrix humilis</i> | 0 | 0 | VU |
| | | <i>Callithrix intermedia</i> | 0 | 0 | LC |
| | | <i>Callithrix kuhlii</i> | 2 | 2 | NT |
| | | <i>Callithrix leucippe</i> | 0 | 0 | |
| | | <i>Callithrix manicorensis</i> | 0 | 0 | |
| | | <i>Callithrix marcai</i> | 0 | 0 | |
| | | <i>Callithrix mauesi</i> | 0 | 0 | |
| | | <i>Callithrix melanura</i> | 0 | 0 | |
| | | <i>Callithrix nigriceps</i> | 0 | 0 | DD |
| | | <i>Callithrix penicillata</i> | 3 | 1 | LC |
| | | <i>Callithrix saterei</i> | 0 | 0 | LC |
| | | <i>Cebus kaapori</i> | 2 | 2 | CR |
| | | <i>Cebus libidinosus</i> | 0 | 0 | LC |
| | | <i>Cebus nigritus</i> | 3 | 3 | NT |
| | | <i>Cebus xanthosternos</i> | 0 | 0 | CR |
| | | <i>Saguinus graellsii</i> | 0 | 0 | |
| | | <i>Saguinus inustus</i> | 2 | 2 | LC |
| | | <i>Saguinus melanoleucus</i> | 0 | 0 | LC |
| | | <i>Saguinus midas midas</i> | 0 | 0 | |
| | | <i>Saguinus niger</i> | 0 | 0 | VU |
| | | <i>Saguinus pileatus</i> | 0 | 0 | |
| | | <i>Saguinus tripartitus</i> | 0 | 0 | NT |
| | | <i>Saimiri ustus</i> | 0 | 0 | NT |
| | | <i>Saimiri vanzolinii</i> | 0 | 0 | VU |
| | CERCOPITHECIDAE | <i>Cercocebus chrysogaster</i> | 0 | 0 | DD |
| | | <i>Cercocebus sanjei</i> | 5 | 5 | EN |
| | | <i>Cercopithecus albogularis</i> | 0 | 0 | LC |
| | | <i>Cercopithecus denti</i> | 0 | 0 | |
| | | <i>Cercopithecus doggetti</i> | 0 | 0 | |
| | | <i>Cercopithecus dryas</i> | 2 | 1 | CR |
| | | <i>Cercopithecus erythrogaster</i> | 0 | 0 | VU |
| | | <i>Cercopithecus lowei</i> | 0 | 0 | |
| | | <i>Cercopithecus preussi</i> | 5 | 5 | EN |
| | | <i>Cercopithecus sclateri</i> | 0 | 0 | VU |
| | | <i>Chlorocebus cynosuros</i> | 0 | 0 | LC |
| | | <i>Chlorocebus djamdjamensis</i> | 0 | 0 | VU |
| | | <i>Lophocebus aterrimus</i> | 2 | 2 | NT |
| | | <i>Lophocebus opdenboschi</i> | 0 | 0 | |
| | | <i>Macaca assamensis</i> | 1 | 1 | NT |
| | | <i>Macaca cyclopis</i> | 3 | 2 | LC |
| | | <i>Macaca hecki</i> | 1 | 1 | VU |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|-----------|-------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Macaca leonina</i> | 0 | 0 | VU |
| | | <i>Macaca maura</i> | 2 | 0 | EN |
| | | <i>Macaca nigrescens</i> | 0 | 0 | VU |
| | | <i>Macaca ochreata</i> | 0 | 0 | VU |
| | | <i>Macaca pagensis</i> | 0 | 0 | CR |
| | | <i>Macaca radiata</i> | 1 | 1 | LC |
| | | <i>Macaca siberu</i> | 0 | 0 | VU |
| | | <i>Macaca thibetana</i> | 0 | 0 | NT |
| | | <i>Macaca tonkeana</i> | 5 | 2 | VU |
| | | <i>Miopithecus ogouensis</i> | 4 | 4 | LC |
| | | <i>Ptilocolobus foai</i> | 3 | 3 | |
| | | <i>Ptilocolobus gordonorum</i> | 0 | 0 | |
| | | <i>Ptilocolobus pennantii</i> | 4 | 4 | |
| | | <i>Ptilocolobus preussi</i> | 0 | 0 | |
| | | <i>Ptilocolobus tephrosceles</i> | 0 | 0 | |
| | | <i>Ptilocolobus tholloni</i> | 0 | 0 | |
| | | <i>Presbytis chrysomelas</i> | 0 | 0 | CR |
| | | <i>Presbytis comata</i> | 4 | 3 | EN |
| | | <i>Presbytis femoralis</i> | 4 | 0 | NT |
| | | <i>Presbytis frontata</i> | 0 | 0 | VU |
| | | <i>Presbytis hosei</i> | 0 | 0 | VU |
| | | <i>Presbytis melalophos</i> | 2 | 2 | EN |
| | | <i>Presbytis natunae</i> | 0 | 0 | VU |
| | | <i>Presbytis rubicunda</i> | 0 | 0 | LC |
| | | <i>Presbytis siamensis</i> | 0 | 0 | NT |
| | | <i>Presbytis thomasi</i> | 0 | 0 | VU |
| | | <i>Trachypithecus barbei</i> | 0 | 0 | DD |
| | | <i>Trachypithecus delacouri</i> | 0 | 0 | CR |
| | | <i>Trachypithecus ebenus</i> | 0 | 0 | EN |
| | | <i>Trachypithecus germaini</i> | 0 | 0 | EN |
| | | <i>Trachypithecus hatinhensis</i> | 0 | 0 | EN |
| | | <i>Trachypithecus laotum</i> | 0 | 0 | VU |
| | | <i>Trachypithecus phayrei</i> | 2 | 1 | EN |
| | | <i>Trachypithecus poliocephalus</i> | 0 | 0 | CR |
| | GALAGIDAE | <i>Euoticus elegantulus</i> | 1 | 1 | LC |
| | | <i>Euoticus pallidus</i> | 0 | 0 | LC |
| | | <i>Galago alleni</i> | 5 | 5 | LC |
| | | <i>Galago cameronensis</i> | 0 | 0 | |
| | | <i>Galago gabonensis</i> | 0 | 0 | |
| | | <i>Galago gallarum</i> | 0 | 0 | LC |
| | | <i>Galago granti</i> | 1 | 1 | LC |
| | | <i>Galago matschiei</i> | 0 | 0 | LC |
| | | <i>Galago moholi</i> | 5 | 5 | LC |
| | | <i>Galago nyasae</i> | 0 | 0 | |
| | | <i>Galago orinus</i> | 1 | 1 | NT |
| | | <i>Galago rondoensis</i> | 0 | 0 | CR |
| | | <i>Galago thomasi</i> | 0 | 0 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|----------|-------------|----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Galago zanzibaricus</i> | 2 | 2 | |
| | | <i>Otolemur garnettii</i> | 5 | 1 | LC |
| | | <i>Otolemur monteiri</i> | 0 | 0 | |
| | LORISIDAE | <i>Arctocebus aureus</i> | 5 | 5 | LC |
| | | <i>Loris lydekkerianus</i> | 0 | 0 | LC |
| | | <i>Pseudopotto martini</i> | 0 | 0 | LC |
| | PITHECIIDAE | <i>Callicebus baptista</i> | 0 | 0 | LC |
| | | <i>Callicebus barbarabrownae</i> | 0 | 0 | CR |
| | | <i>Callicebus bernhardi</i> | 0 | 0 | LC |
| | | <i>Callicebus brunneus</i> | 4 | 4 | LC |
| | | <i>Callicebus caligatus</i> | 0 | 0 | LC |
| | | <i>Callicebus cinerascens</i> | 0 | 0 | LC |
| | | <i>Callicebus coimbrai</i> | 0 | 0 | EN |
| | | <i>Callicebus donacophilus</i> | 0 | 0 | LC |
| | | <i>Callicebus dubius</i> | 0 | 0 | LC |
| | | <i>Callicebus hoffmannsi</i> | 0 | 0 | LC |
| | | <i>Callicebus lucifer</i> | 0 | 0 | LC |
| | | <i>Callicebus lugens</i> | 0 | 0 | LC |
| | | <i>Callicebus medemi</i> | 0 | 0 | VU |
| | | <i>Callicebus melanochir</i> | 0 | 0 | VU |
| | | <i>Callicebus modestus</i> | 0 | 0 | EN |
| | | <i>Callicebus nigrifrons</i> | 0 | 0 | NT |
| | | <i>Callicebus oenanthe</i> | 0 | 0 | EN |
| | | <i>Callicebus olallae</i> | 0 | 0 | EN |
| | | <i>Callicebus ornatus</i> | 0 | 0 | VU |
| | | <i>Callicebus pallescens</i> | 0 | 0 | LC |
| | | <i>Callicebus personatus</i> | 0 | 0 | VU |
| | | <i>Callicebus purinus</i> | 0 | 0 | LC |
| | | <i>Callicebus regulus</i> | 0 | 0 | LC |
| | | <i>Callicebus stephennashi</i> | 0 | 0 | DD |
| | | <i>Callicebus torquatus</i> | 0 | 0 | LC |
| | | <i>Chiropotes chiropotes</i> | 5 | 5 | LC |
| | | <i>Chiropotes israelita</i> | 0 | 0 | LC |
| | | <i>Chiropotes utahickae</i> | 0 | 0 | EN |
| | | <i>Pithecia albicans</i> | 0 | 0 | VU |
| | | <i>Pithecia irrorata</i> | 2 | 2 | LC |
| | TARSIIDAE | <i>Tarsius bancanus</i> | 1 | 1 | VU |
| | | <i>Tarsius lariang</i> | 2 | 2 | DD |
| | | <i>Tarsius pelengensis</i> | 0 | 0 | EN |
| | | <i>Tarsius pumilus</i> | 1 | 1 | DD |
| | | <i>Tarsius sangirensis</i> | 0 | 0 | EN |
| | | <i>Tarsius syrichta</i> | 5 | 5 | NT |
| RODENTIA | SCIURIDAE | <i>Ratufa affinis</i> | 1 | 1 | NT |
| | | <i>Ratufa bicolor</i> | 2 | 1 | NT |
| | | <i>Ratufa indica</i> | 0 | 0 | LC |
| | | <i>Ratufa macroura</i> | 0 | 0 | NT |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------------------|---------------|--------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| SCANDENTIA | PTILOCERCIDAE | <i>Ptilocercus lowii</i> | 0 | 0 | LC |
| | TUPAIIDAE | <i>Anathana ellioti</i> | 0 | 0 | LC |
| | | <i>Dendrogale melanura</i> | 0 | 0 | DD |
| | | <i>Dendrogale murina</i> | 0 | 0 | LC |
| | | <i>Tupaia chrysogaster</i> | 0 | 0 | EN |
| | | <i>Tupaia dorsalis</i> | 1 | 1 | DD |
| | | <i>Tupaia gracilis</i> | 0 | 0 | LC |
| | | <i>Tupaia javanica</i> | 4 | 4 | LC |
| | | <i>Tupaia longipes</i> | 0 | 0 | LC |
| | | <i>Tupaia moellendorffi</i> | 0 | 0 | DD |
| | | <i>Tupaia nicobarica</i> | 0 | 0 | EN |
| | | <i>Tupaia picta</i> | 1 | 1 | LC |
| | | <i>Tupaia splendidula</i> | 0 | 0 | LC |
| | | <i>Tupaia tana</i> | 2 | 2 | LC |
| | | <i>Urogale everetti</i> | 1 | 1 | LC |
| CLASS AVES | | | | | |
| ANSERIFORMES | ANATIDAE | <i>Dendrocygna arborea</i> | 4 | 3 | VU |
| | | <i>Oxyura leucocephala</i> | 3 | 2 | EN |
| APODIFORMES | TROCHILIDAE | <i>Abeillia abeillei</i> | 2 | 2 | LC |
| | | <i>Aglaeactis aliciae</i> | 0 | 0 | EN |
| | | <i>Aglaeactis castelnaudii</i> | 3 | 3 | LC |
| | | <i>Aglaeactis pamela</i> | 0 | 0 | LC |
| | | <i>Aglaiocercus coelestis</i> | 1 | 0 | LC |
| | | <i>Amazilia alticola</i> | 0 | 0 | |
| | | <i>Amazilia boucardi</i> | 1 | 0 | EN |
| | | <i>Amazilia castaneiventris</i> | 1 | 0 | EN |
| | | <i>Amazilia cupreicauda</i> | 1 | 1 | |
| | | <i>Amazilia cyanifrons</i> | 1 | 0 | LC |
| | | <i>Amazilia cyanura</i> | 5 | 3 | LC |
| | | <i>Amazilia edward</i> | 4 | 2 | LC |
| | | <i>Amazilia luciae</i> | 0 | 0 | CR |
| | | <i>Amazilia rondoniae</i> | 0 | 0 | |
| | | <i>Amazilia rosenbergi</i> | 3 | 2 | LC |
| | | <i>Amazilia saucerrottei</i> | 0 | 0 | LC |
| | | <i>Amazilia tobaci</i> | 5 | 4 | LC |
| | | <i>Amazilia violiceps</i> | 3 | 3 | LC |
| | | <i>Amazilia viridifrons</i> | 5 | 2 | LC |
| | | <i>Androdon aequatorialis</i> | 1 | 0 | LC |
| | | <i>Anopetia gounellei</i> | 1 | 1 | LC |
| | | <i>Anthocephala floriceps</i> | 0 | 0 | VU |
| | | <i>Anthracothorax dominicus</i> | 2 | 2 | LC |
| | | <i>Anthracothorax recurvirostris</i> | 2 | 2 | |
| | | <i>Anthracothorax veraguensis</i> | 2 | 2 | LC |
| | | <i>Anthracothorax viridigula</i> | 4 | 2 | LC |
| | | <i>Anthracothorax viridis</i> | 1 | 1 | LC |
| | | <i>Aphantochroa cirrochloris</i> | 0 | 0 | LC |
| | | <i>Atthis ellioti</i> | 0 | 0 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|--------|-------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Augastes lumachella</i> | 0 | 0 | NT |
| | | <i>Augastes scutatus</i> | 2 | 0 | NT |
| | | <i>Basilinna xantusii</i> | 3 | 2 | LC |
| | | <i>Boissonneaua flavescens</i> | 3 | 2 | LC |
| | | <i>Boissonneaua jardini</i> | 2 | 0 | LC |
| | | <i>Calliphlox bryantae</i> | 2 | 0 | LC |
| | | <i>Calliphlox mitchellii</i> | 0 | 0 | LC |
| | | <i>Calothorax lucifer</i> | 1 | 0 | LC |
| | | <i>Calothorax pulcher</i> | 2 | 1 | LC |
| | | <i>Campylopterus curvipennis</i> | 4 | 4 | LC |
| | | <i>Campylopterus cuvierii</i> | 5 | 5 | |
| | | <i>Campylopterus duidae</i> | 1 | 0 | LC |
| | | <i>Campylopterus falcatus</i> | 4 | 4 | LC |
| | | <i>Campylopterus hyperythrus</i> | 3 | 3 | LC |
| | | <i>Campylopterus phainopeplus</i> | 2 | 0 | EN |
| | | <i>Campylopterus villaviscensio</i> | 4 | 3 | NT |
| | | <i>Chaetocercus astreans</i> | 0 | 0 | LC |
| | | <i>Chaetocercus berlepschi</i> | 0 | 0 | EN |
| | | <i>Chaetocercus heliodor</i> | 4 | 4 | LC |
| | | <i>Chaetocercus jourdani</i> | 0 | 0 | LC |
| | | <i>Chalcostigma heteropogon</i> | 2 | 1 | LC |
| | | <i>Chalcostigma stanleyi</i> | 2 | 2 | LC |
| | | <i>Chlorostilbon alice</i> | 0 | 0 | LC |
| | | <i>Chlorostilbon assimilis</i> | 4 | 4 | LC |
| | | <i>Chlorostilbon auriceps</i> | 0 | 0 | LC |
| | | <i>Chlorostilbon forficatus</i> | 0 | 0 | LC |
| | | <i>Chlorostilbon maugaeus</i> | 1 | 1 | LC |
| | | <i>Chlorostilbon olivaresi</i> | 0 | 0 | LC |
| | | <i>Chlorostilbon poortmani</i> | 0 | 0 | LC |
| | | <i>Chlorostilbon ricordii</i> | 3 | 3 | LC |
| | | <i>Chlorostilbon russatus</i> | 1 | 1 | LC |
| | | <i>Chlorostilbon stenurus</i> | 3 | 2 | LC |
| | | <i>Chlorostilbon swainsonii</i> | 3 | 3 | LC |
| | | <i>Clytolaema rubricauda</i> | 4 | 3 | LC |
| | | <i>Coeligena bonapartei</i> | 0 | 0 | LC |
| | | <i>Coeligena helianthea</i> | 1 | 0 | LC |
| | | <i>Coeligena lutetiae</i> | 2 | 2 | LC |
| | | <i>Coeligena phalerata</i> | 0 | 0 | LC |
| | | <i>Coeligena prunellei</i> | 0 | 0 | VU |
| | | <i>Coeligena wilsoni</i> | 0 | 0 | LC |
| | | <i>Colibri serrirostris</i> | 3 | 1 | LC |
| | | <i>Cyanophaia bicolor</i> | 0 | 0 | LC |
| | | <i>Cynanthus sordidus</i> | 2 | 1 | LC |
| | | <i>Discosura conversii</i> | 0 | 0 | LC |
| | | <i>Discosura langsdorffi</i> | 2 | 1 | LC |
| | | <i>Discosura letitiae</i> | 0 | 0 | DD |
| | | <i>Discosura longicaudus</i> | 1 | 1 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|--------|-----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Discosura popelairii</i> | 1 | 0 | LC |
| | | <i>Doricha eliza</i> | 2 | 2 | NT |
| | | <i>Doricha enicura</i> | 0 | 0 | LC |
| | | <i>Elvira chionura</i> | 3 | 2 | LC |
| | | <i>Elvira cupreiceps</i> | 3 | 0 | LC |
| | | <i>Eriocnemis cupreiventris</i> | 0 | 0 | NT |
| | | <i>Eriocnemis derbyi</i> | 2 | 1 | NT |
| | | <i>Eriocnemis godini</i> | 0 | 0 | CR |
| | | <i>Eriocnemis isabellae</i> | 0 | 0 | CR |
| | | <i>Eriocnemis mirabilis</i> | 0 | 0 | CR |
| | | <i>Eriocnemis mosquera</i> | 1 | 0 | LC |
| | | <i>Eriocnemis nigrivestis</i> | 5 | 1 | CR |
| | | <i>Eriocnemis vestita</i> | 1 | 1 | LC |
| | | <i>Eulampis holosericeus</i> | 1 | 1 | LC |
| | | <i>Eulampis jugularis</i> | 3 | 2 | LC |
| | | <i>Eulidia yarrellii</i> | 0 | 0 | EN |
| | | <i>Eupetomena macroura</i> | 0 | 0 | LC |
| | | <i>Eupherusa cyanophrys</i> | 3 | 3 | EN |
| | | <i>Eupherusa nigriventris</i> | 2 | 2 | LC |
| | | <i>Eupherusa poliocerca</i> | 1 | 1 | VU |
| | | <i>Glaucis aeneus</i> | 4 | 4 | LC |
| | | <i>Goethalsia bella</i> | 0 | 0 | NT |
| | | <i>Goldmania violiceps</i> | 2 | 2 | LC |
| | | <i>Haplophaedia assimilis</i> | 5 | 4 | LC |
| | | <i>Haplophaedia lugens</i> | 0 | 0 | NT |
| | | <i>Heliangelus exortis</i> | 3 | 3 | LC |
| | | <i>Heliangelus mavors</i> | 0 | 0 | LC |
| | | <i>Heliangelus regalis</i> | 4 | 4 | EN |
| | | <i>Heliangelus strophianus</i> | 1 | 0 | LC |
| | | <i>Heliodoxa gularis</i> | 3 | 3 | NT |
| | | <i>Heliodoxa imperatrix</i> | 1 | 1 | LC |
| | | <i>Heliodoxa schreibersii</i> | 2 | 1 | LC |
| | | <i>Heliomaster constantii</i> | 3 | 2 | LC |
| | | <i>Heliothyx barroti</i> | 4 | 3 | LC |
| | | <i>Hylocharis grayi</i> | 1 | 0 | LC |
| | | <i>Hylonympha macrocerca</i> | 0 | 0 | EN |
| | | <i>Lampornis calolaemus</i> | 0 | 0 | LC |
| | | <i>Lampornis hemileucus</i> | 4 | 4 | LC |
| | | <i>Lampornis sybillae</i> | 0 | 0 | LC |
| | | <i>Lamprolaima rhami</i> | 2 | 2 | LC |
| | | <i>Lepidopyga coeruleogularis</i> | 5 | 5 | LC |
| | | <i>Lepidopyga goudoti</i> | 0 | 0 | LC |
| | | <i>Lepidopyga lilliae</i> | 0 | 0 | CR |
| | | <i>Leucippus baeri</i> | 1 | 0 | LC |
| | | <i>Leucippus chlorocercus</i> | 3 | 2 | LC |
| | | <i>Leucippus fallax</i> | 1 | 0 | LC |
| | | <i>Loddigesia mirabilis</i> | 0 | 0 | EN |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|--------|-----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Lophornis adorabilis</i> | 0 | 0 | LC |
| | | <i>Lophornis brachylophus</i> | 0 | 0 | CR |
| | | <i>Lophornis chalybeus</i> | 2 | 0 | LC |
| | | <i>Lophornis delattrei</i> | 2 | 2 | LC |
| | | <i>Lophornis gouldii</i> | 0 | 0 | LC |
| | | <i>Lophornis helenae</i> | 1 | 0 | LC |
| | | <i>Lophornis stictolophus</i> | 3 | 3 | LC |
| | | <i>Mellisuga helenae</i> | 2 | 0 | NT |
| | | <i>Mellisuga minima</i> | 5 | 4 | LC |
| | | <i>Metallura baroni</i> | 1 | 1 | EN |
| | | <i>Metallura iracunda</i> | 0 | 0 | EN |
| | | <i>Metallura odomae</i> | 0 | 0 | NT |
| | | <i>Metallura theresiae</i> | 3 | 3 | LC |
| | | <i>Metallura williami</i> | 2 | 2 | LC |
| | | <i>Microchera albocoronata</i> | 3 | 3 | LC |
| | | <i>Microstilbon burmeisteri</i> | 1 | 1 | LC |
| | | <i>Myrmia micrura</i> | 0 | 0 | LC |
| | | <i>Opisthoprora euryptera</i> | 2 | 2 | LC |
| | | <i>Oreonympha nobilis</i> | 4 | 4 | LC |
| | | <i>Oreotrochilus adela</i> | 0 | 0 | NT |
| | | <i>Oreotrochilus chimborazo</i> | 3 | 1 | LC |
| | | <i>Orthorhyncus cristatus</i> | 1 | 1 | LC |
| | | <i>Oxypogon guerinii</i> | 1 | 1 | LC |
| | | <i>Panterpe insignis</i> | 0 | 0 | LC |
| | | <i>Phaethornis anthophilus</i> | 2 | 2 | LC |
| | | <i>Phaethornis atrimentalis</i> | 2 | 2 | LC |
| | | <i>Phaethornis eurynome</i> | 5 | 4 | LC |
| | | <i>Phaethornis idaliae</i> | 1 | 1 | LC |
| | | <i>Phaethornis koepckeae</i> | 5 | 5 | NT |
| | | <i>Phaethornis longirostris</i> | 5 | 5 | LC |
| | | <i>Phaethornis nattereri</i> | 0 | 0 | LC |
| | | <i>Phaethornis philippii</i> | 1 | 1 | LC |
| | | <i>Phaethornis pretrei</i> | 0 | 0 | LC |
| | | <i>Phaethornis rupurumii</i> | 2 | 2 | LC |
| | | <i>Phaethornis squalidus</i> | 1 | 0 | LC |
| | | <i>Phaethornis stuarti</i> | 5 | 5 | LC |
| | | <i>Phaethornis subochraceus</i> | 0 | 0 | LC |
| | | <i>Phaethornis yaruqui</i> | 1 | 1 | LC |
| | | <i>Phlogophilus harterti</i> | 4 | 4 | NT |
| | | <i>Phlogophilus hemileucurus</i> | 0 | 0 | NT |
| | | <i>Polytmus milleri</i> | 2 | 1 | LC |
| | | <i>Ramphodon naevius</i> | 1 | 1 | NT |
| | | <i>Ramphomicron dorsale</i> | 0 | 0 | LC |
| | | <i>Ramphomicron microrhynchum</i> | 1 | 1 | LC |
| | | <i>Selasphorus ardens</i> | 3 | 3 | VU |
| | | <i>Selasphorus sasin</i> | 3 | 2 | LC |
| | | <i>Selasphorus scintilla</i> | 3 | 0 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|---------------|------------------|-----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Stellula calliope</i> | 3 | 3 | LC |
| | | <i>Stephanoxis lalandi</i> | 5 | 4 | LC |
| | | <i>Sternoclyta cyanopectus</i> | 1 | 0 | LC |
| | | <i>Taphrolesia griseiventris</i> | 0 | 0 | EN |
| | | <i>Thalurania fannyi</i> | 3 | 2 | LC |
| | | <i>Thalurania glaucopis</i> | 3 | 3 | LC |
| | | <i>Thalurania ridgwayi</i> | 1 | 1 | VU |
| | | <i>Thalurania watertonii</i> | 0 | 0 | NT |
| | | <i>Tilmatura dupontii</i> | 1 | 1 | LC |
| | | <i>Trochilus scitulus</i> | 0 | 0 | LC |
| | | <i>Urochroa bougueri</i> | 5 | 3 | LC |
| | | <i>Urosticte benjamini</i> | 2 | 2 | LC |
| CICONIIFORMES | PHOENICOPTERIDAE | <i>Phoenicopterus ruber ruber</i> | 0 | 0 | LC |
| COLUMBIFORMES | COLUMBIDAE | <i>Goura cristata</i> | 2 | 0 | VU |
| CORACIIFORMES | BUCEROTIDAE | <i>Aceros cassidix</i> | 4 | 0 | LC |
| | | <i>Aceros leucocephalus</i> | 1 | 1 | NT |
| | | <i>Aceros waldeni</i> | 3 | 3 | CR |
| | | <i>Anorrhinus galeritus</i> | 1 | 0 | LC |
| | | <i>Anthracoceros malayanus</i> | 3 | 1 | NT |
| | | <i>Anthracoceros marchei</i> | 0 | 0 | VU |
| | | <i>Anthracoceros montani</i> | 0 | 0 | CR |
| | | <i>Bericornis comatus</i> | 2 | 0 | |
| | | <i>Rhyticeros everetti</i> | 0 | 0 | VU |
| | | <i>Rhyticeros narcondami</i> | 2 | 1 | |
| CUCULIFORMES | MUSOPHAGIDAE | <i>Tauraco bannermani</i> | 2 | 0 | EN |
| | | <i>Tauraco ruspolii</i> | 0 | 0 | VU |
| | | <i>Tauraco schuettii</i> | 3 | 1 | LC |
| FALCONIFORMES | ACCIPITRIDAE | <i>Accipiter albobularis</i> | 1 | 1 | LC |
| | | <i>Accipiter brachyurus</i> | 0 | 0 | VU |
| | | <i>Accipiter brevipes</i> | 4 | 4 | LC |
| | | <i>Accipiter butleri</i> | 2 | 1 | VU |
| | | <i>Accipiter castanilius</i> | 3 | 3 | LC |
| | | <i>Accipiter cirrocephalus</i> | 0 | 0 | LC |
| | | <i>Accipiter collaris</i> | 1 | 1 | NT |
| | | <i>Accipiter erythrauchen</i> | 0 | 0 | LC |
| | | <i>Accipiter erythropus</i> | 4 | 4 | LC |
| | | <i>Accipiter francesii</i> | 0 | 0 | LC |
| | | <i>Accipiter griseiceps</i> | 0 | 0 | LC |
| | | <i>Accipiter gundlachi</i> | 0 | 0 | EN |
| | | <i>Accipiter haplochrous</i> | 1 | 1 | NT |
| | | <i>Accipiter henicogrammus</i> | 0 | 0 | LC |
| | | <i>Accipiter henstii</i> | 3 | 3 | NT |
| | | <i>Accipiter imitator</i> | 1 | 1 | VU |
| | | <i>Accipiter luteoschistaceus</i> | 0 | 0 | VU |
| | | <i>Accipiter madagascariensis</i> | 2 | 1 | NT |
| | | <i>Accipiter melanochlamys</i> | 3 | 3 | LC |
| | | <i>Accipiter meyerianus</i> | 0 | 0 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|--------|-----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Accipiter nanus</i> | 0 | 0 | NT |
| | | <i>Accipiter ovampensis</i> | 2 | 2 | LC |
| | | <i>Accipiter poliogaster</i> | 4 | 4 | LC |
| | | <i>Accipiter princeps</i> | 0 | 0 | VU |
| | | <i>Accipiter rhodogaster</i> | 0 | 0 | LC |
| | | <i>Accipiter rufitorques</i> | 0 | 0 | LC |
| | | <i>Accipiter toussenelii</i> | 2 | 2 | |
| | | <i>Accipiter trinotatus</i> | 0 | 0 | LC |
| | | <i>Accipiter trivirgatus</i> | 4 | 4 | LC |
| | | <i>Aquila gurneyi</i> | 0 | 0 | NT |
| | | <i>Aquila hastata</i> | 0 | 0 | VU |
| | | <i>Aviceda jerdoni</i> | 2 | 2 | LC |
| | | <i>Aviceda leuphotes</i> | 2 | 2 | LC |
| | | <i>Aviceda madagascariensis</i> | 2 | 2 | LC |
| | | <i>Butastur indicus</i> | 2 | 1 | LC |
| | | <i>Butastur liventer</i> | 0 | 0 | LC |
| | | <i>Butastur teesa</i> | 0 | 0 | LC |
| | | <i>Buteo albigula</i> | 2 | 2 | LC |
| | | <i>Buteo augur</i> | 4 | 4 | LC |
| | | <i>Buteo brachypterus</i> | 3 | 3 | LC |
| | | <i>Buteo leucorrhous</i> | 5 | 5 | LC |
| | | <i>Buteo solitarius</i> | 0 | 0 | NT |
| | | <i>Buteo ventralis</i> | 0 | 0 | NT |
| | | <i>Buteogallus anthracinus</i> | 4 | 3 | LC |
| | | <i>Buteogallus subtilis</i> | 2 | 2 | |
| | | <i>Chelictinia riocourii</i> | 1 | 1 | LC |
| | | <i>Chondrohierax uncinatus</i> | 4 | 4 | LC |
| | | <i>Circaetus beaudouini</i> | 0 | 0 | VU |
| | | <i>Circaetus cinerascens</i> | 4 | 3 | LC |
| | | <i>Circaetus fasciolatus</i> | 1 | 1 | NT |
| | | <i>Circus assimilis</i> | 1 | 1 | LC |
| | | <i>Circus cinereus</i> | 3 | 2 | LC |
| | | <i>Circus maillardi</i> | 0 | 0 | EN |
| | | <i>Circus maurus</i> | 1 | 1 | VU |
| | | <i>Circus melanoleucos</i> | 1 | 1 | LC |
| | | <i>Circus pygargus</i> | 2 | 2 | LC |
| | | <i>Circus ranivorus</i> | 2 | 1 | LC |
| | | <i>Circus spilonotus</i> | 1 | 0 | LC |
| | | <i>Dryotriorchis spectabilis</i> | 1 | 1 | LC |
| | | <i>Elanus axillaris</i> | 5 | 4 | LC |
| | | <i>Elanus leucurus</i> | 3 | 3 | LC |
| | | <i>Elanus scriptus</i> | 2 | 2 | NT |
| | | <i>Erythrotriorchis buergersi</i> | 1 | 1 | DD |
| | | <i>Erythrotriorchis radiatus</i> | 1 | 1 | VU |
| | | <i>Eutriorchis astur</i> | 4 | 4 | EN |
| | | <i>Gyps indicus</i> | 4 | 4 | CR |
| | | <i>Gyps tenuirostris</i> | 2 | 2 | CR |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|------------|----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Haliaeetus leucoryphus</i> | 1 | 0 | VU |
| | | <i>Haliaeetus sanfordi</i> | 1 | 1 | VU |
| | | <i>Haliastur sphenurus</i> | 1 | 1 | LC |
| | | <i>Hamirostra melanosternon</i> | 1 | 1 | LC |
| | | <i>Harpagus diodon</i> | 0 | 0 | LC |
| | | <i>Harpyhaliaetus coronatus</i> | 5 | 3 | EN |
| | | <i>Harpyhaliaetus solitarius</i> | 2 | 2 | NT |
| | | <i>Harpyopsis novaeguineae</i> | 4 | 4 | VU |
| | | <i>Henicopernis infuscatus</i> | 1 | 0 | VU |
| | | <i>Hieraaetus kienerii</i> | 1 | 1 | LC |
| | | <i>Hieraaetus morphnoides</i> | 3 | 3 | LC |
| | | <i>Ichthyophaga humilis</i> | 1 | 0 | NT |
| | | <i>Ichthyophaga ichthyaetus</i> | 1 | 1 | NT |
| | | <i>Ictinaetus malayensis</i> | 2 | 1 | LC |
| | | <i>Ictinia mississippiensis</i> | 2 | 2 | LC |
| | | <i>Leptodon cayanensis</i> | 3 | 3 | LC |
| | | <i>Leucopternis kuhli</i> | 4 | 4 | LC |
| | | <i>Leucopternis lacernulatus</i> | 0 | 0 | VU |
| | | <i>Leucopternis occidentalis</i> | 2 | 2 | EN |
| | | <i>Leucopternis plumbeus</i> | 2 | 2 | NT |
| | | <i>Leucopternis polionotus</i> | 0 | 0 | NT |
| | | <i>Leucopternis princeps</i> | 1 | 1 | LC |
| | | <i>Leucopternis schistaceus</i> | 4 | 4 | LC |
| | | <i>Leucopternis semiplumbeus</i> | 3 | 3 | LC |
| | | <i>Lophoictinia isura</i> | 1 | 1 | LC |
| | | <i>Macheiramphus alcinus</i> | 4 | 4 | LC |
| | | <i>Megatriorchis doriae</i> | 4 | 4 | NT |
| | | <i>Morphnus guianensis</i> | 4 | 4 | NT |
| | | <i>Oroaetus isidori</i> | 2 | 2 | VU |
| | | <i>Pernis celebensis</i> | 1 | 1 | LC |
| | | <i>Polyboroides radiatus</i> | 3 | 3 | LC |
| | | <i>Rostrhamus hamatus</i> | 0 | 0 | LC |
| | | <i>Spilornis elgini</i> | 0 | 0 | NT |
| | | <i>Spilornis kinabaluensis</i> | 1 | 1 | VU |
| | | <i>Spilornis klossi</i> | 0 | 0 | NT |
| | | <i>Spilornis rufipectus</i> | 0 | 0 | LC |
| | | <i>Spizaetus africanus</i> | 4 | 4 | LC |
| | | <i>Spizaetus alboniger</i> | 2 | 2 | LC |
| | | <i>Spizaetus bartelsi</i> | 0 | 0 | EN |
| | | <i>Spizaetus lanceolatus</i> | 0 | 0 | LC |
| | | <i>Spizaetus nanus</i> | 2 | 2 | VU |
| | | <i>Spizaetus philippensis</i> | 1 | 1 | VU |
| | | <i>Spizastur melanoleucus</i> | 5 | 4 | LC |
| | | <i>Urotriorchis macrourus</i> | 5 | 5 | LC |
| | FALCONIDAE | <i>Caracara cheriway</i> | 1 | 1 | LC |
| | | <i>Caracara lutosa</i> | 0 | 0 | EX |
| | | <i>Falco berigora</i> | 5 | 5 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|---------------|---------------|----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Falco concolor</i> | 2 | 2 | NT |
| | | <i>Falco dickinsoni</i> | 0 | 0 | LC |
| | | <i>Falco fasciinucha</i> | 5 | 0 | NT |
| | | <i>Falco hypoleucos</i> | 2 | 1 | NT |
| | | <i>Falco longipennis</i> | 2 | 2 | LC |
| | | <i>Falco moluccensis</i> | 0 | 0 | LC |
| | | <i>Falco newtoni</i> | 3 | 3 | LC |
| | | <i>Falco severus</i> | 4 | 4 | LC |
| | | <i>Falco subniger</i> | 1 | 1 | LC |
| | | <i>Falco zoniventris</i> | 1 | 1 | LC |
| | | <i>Micrastur buckleyi</i> | 4 | 4 | LC |
| | | <i>Micrastur mintoni</i> | 0 | 0 | LC |
| | | <i>Micrastur mirandollei</i> | 5 | 5 | LC |
| | | <i>Micrastur plumbeus</i> | 0 | 0 | VU |
| | | <i>Microhierax erythrogenys</i> | 2 | 2 | LC |
| | | <i>Microhierax fringillarius</i> | 0 | 0 | LC |
| | | <i>Microhierax latifrons</i> | 0 | 0 | NT |
| | | <i>Microhierax melanoleucos</i> | 0 | 0 | LC |
| | | <i>Phalcoboenus albogularis</i> | 1 | 1 | LC |
| | | <i>Phalcoboenus carunculatus</i> | 0 | 0 | LC |
| | | <i>Polihierax insignis</i> | 0 | 0 | NT |
| | | <i>Spiziapteryx circumcincta</i> | 5 | 5 | LC |
| GRUIFORMES | GRUIDAE | <i>Grus rubicunda</i> | 3 | 0 | LC |
| | OTIDIDAE | <i>Ardeotis australis</i> | 3 | 3 | NT |
| | | <i>Eupodotis caerulescens</i> | 3 | 3 | NT |
| | | <i>Eupodotis humilis</i> | 4 | 3 | NT |
| | | <i>Lissotis hartlaubii</i> | 3 | 3 | |
| | | <i>Lophotis savilei</i> | 2 | 2 | |
| | | <i>Neotis heuglinii</i> | 4 | 4 | LC |
| | | <i>Neotis ludwigii</i> | 5 | 5 | EN |
| | | <i>Sypheotides indicus</i> | 2 | 2 | EN |
| PASSERIFORMES | COTINGIDAE | <i>Rupicola peruvianus</i> | 0 | 0 | LC |
| | EMBERIZIDAE | <i>Gubernatrix cristata</i> | 2 | 2 | EN |
| | | <i>Tangara fastuosa</i> | 3 | 3 | VU |
| | ESTRILDIDAE | <i>Amandava formosa</i> | 1 | 0 | VU |
| | | <i>Poephila cincta cincta</i> | 0 | 0 | NT |
| | FRINGILLIDAE | <i>Carduelis yarrellii</i> | 0 | 0 | VU |
| | MUSCICAPIDAE | <i>Cyornis ruckii</i> | 0 | 0 | CR |
| | | <i>Garrulax canorus</i> | 3 | 1 | LC |
| | | <i>Garrulax taewanus</i> | 0 | 0 | NT |
| | | <i>Liocichla omeiensis</i> | 2 | 1 | VU |
| | PARADISAEIDAE | <i>Astrapia mayeri</i> | 2 | 2 | NT |
| | | <i>Astrapia nigra</i> | 0 | 0 | LC |
| | | <i>Astrapia rothschildi</i> | 0 | 0 | LC |
| | | <i>Astrapia splendidissima</i> | 0 | 0 | LC |
| | | <i>Astrapia stephaniae</i> | 5 | 5 | LC |
| | | <i>Cnemophilus loriae</i> | 3 | 3 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|----------------|------------|------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Cnemophilus macgregorii</i> | 0 | 0 | LC |
| | | <i>Diphylloides respublica</i> | 0 | 0 | NT |
| | | <i>Drepanornis albertisi</i> | 1 | 1 | |
| | | <i>Drepanornis bruijnii</i> | 0 | 0 | NT |
| | | <i>Epimachus meyeri</i> | 2 | 2 | LC |
| | | <i>Loboparadisea sericea</i> | 5 | 5 | NT |
| | | <i>Lycocorax pyrrhopterus</i> | 0 | 0 | LC |
| | | <i>Macgregoria pulchra</i> | 0 | 0 | VU |
| | | <i>Manucodia ater</i> | 5 | 4 | LC |
| | | <i>Manucodia chalybatus</i> | 0 | 0 | LC |
| | | <i>Manucodia comrii</i> | 0 | 0 | LC |
| | | <i>Manucodia jobiensis</i> | 0 | 0 | LC |
| | | <i>Melampitta gigantea</i> | 0 | 0 | LC |
| | | <i>Melampitta lugubris</i> | 0 | 0 | LC |
| | | <i>Paradigalla brevicauda</i> | 0 | 0 | LC |
| | | <i>Paradigalla carunculata</i> | 0 | 0 | NT |
| | | <i>Paradisaea decora</i> | 1 | 1 | NT |
| | | <i>Paradisaea guilielmi</i> | 1 | 1 | NT |
| | | <i>Parotia carolae</i> | 3 | 3 | LC |
| | | <i>Parotia helenae</i> | 0 | 0 | LC |
| | | <i>Parotia lawesii</i> | 0 | 0 | LC |
| | | <i>Parotia sefilata</i> | 0 | 0 | LC |
| | | <i>Parotia wahnesi</i> | 0 | 0 | VU |
| | | <i>Phonygammus keraudrenii</i> | 2 | 2 | |
| | | <i>Ptiloris intercedens</i> | 0 | 0 | LC |
| | | <i>Ptiloris paradiseus</i> | 4 | 0 | LC |
| | | <i>Ptiloris victoriae</i> | 5 | 2 | LC |
| | | <i>Semioptera wallacii</i> | 2 | 0 | LC |
| | PITTIDAE | <i>Pitta nympha</i> | 2 | 0 | VU |
| PSITTACIFORMES | CACATUIDAE | <i>Cacatua pastinator</i> | 3 | 0 | LC |
| | | <i>Calyptorhynchus baudinii</i> | 3 | 0 | EN |
| | | <i>Calyptorhynchus lathamii</i> | 5 | 3 | LC |
| | | <i>Calyptorhynchus latirostris</i> | 0 | 0 | EN |
| | LORIIDAE | <i>Chamosyna aureicincta</i> | 0 | 0 | |
| | | <i>Chamosyna diadema</i> | 0 | 0 | CR |
| | | <i>Chamosyna josefinae</i> | 5 | 5 | LC |
| | | <i>Chamosyna margarethae</i> | 4 | 4 | NT |
| | | <i>Chamosyna meeki</i> | 1 | 1 | NT |
| | | <i>Chamosyna multistriata</i> | 3 | 3 | NT |
| | | <i>Chamosyna palmarum</i> | 0 | 0 | VU |
| | | <i>Chamosyna placentis ornata</i> | 0 | 0 | LC |
| | | <i>Chamosyna rubrigularis</i> | 0 | 0 | LC |
| | | <i>Chamosyna rubronotata</i> | 2 | 2 | LC |
| | | <i>Chamosyna toxopei</i> | 0 | 0 | CR |
| | | <i>Chamosyna wilhelminae</i> | 2 | 2 | LC |
| | | <i>Eos cyanogenia</i> | 1 | 1 | VU |
| | | <i>Eos semilarvata</i> | 3 | 0 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|-------------|--------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Eos squamata riciniata</i> | 0 | 0 | |
| | | <i>Glossopsitta porphyrocephala</i> | 5 | 5 | LC |
| | | <i>Lorius albidinucha</i> | 4 | 1 | NT |
| | | <i>Neopsittacus pullicauda</i> | 0 | 0 | LC |
| | | <i>Psitteuteles iris</i> | 0 | 0 | NT |
| | | <i>Trichoglossus euteles</i> | 5 | 0 | LC |
| | | <i>Trichoglossus johnstoniae</i> | 5 | 3 | NT |
| | | <i>Trichoglossus rubiginosus</i> | 0 | 0 | LC |
| | | <i>Vini kuhlii</i> | 5 | 5 | EN |
| | | <i>Vini peruviana</i> | 2 | 0 | VU |
| | | <i>Vini stepheni</i> | 1 | 1 | VU |
| | PSITTACIDAE | <i>Amazona agilis</i> | 4 | 2 | VU |
| | | <i>Amazona festiva festiva</i> | 0 | 0 | LC |
| | | <i>Amazona kawalli</i> | 2 | 2 | LC |
| | | <i>Aprosmictus jonquillaceus</i> | 3 | 0 | NT |
| | | <i>Aratinga cactorum</i> | 0 | 0 | LC |
| | | <i>Aratinga euops</i> | 5 | 1 | VU |
| | | <i>Aratinga maculata</i> | 0 | 0 | |
| | | <i>Bolborhynchus ferrugineifrons</i> | 0 | 0 | VU |
| | | <i>Coracopsis nigra barklyi</i> | 0 | 0 | LC |
| | | <i>Cyanoramphus malherbi</i> | 0 | 0 | CR |
| | | <i>Cyanoramphus unicolor</i> | 2 | 0 | VU |
| | | <i>Enicognathus ferrugineus</i> | 4 | 1 | LC |
| | | <i>Enicognathus leptorhynchus</i> | 5 | 2 | LC |
| | | <i>Forpus xanthops</i> | 4 | 3 | VU |
| | | <i>Geoffroyus simplex</i> | 0 | 0 | LC |
| | | <i>Hapalopsittaca amazonina</i> | 2 | 2 | VU |
| | | <i>Hapalopsittaca fuertesi</i> | 0 | 0 | CR |
| | | <i>Hapalopsittaca melanotis</i> | 4 | 4 | LC |
| | | <i>Hapalopsittaca pyrrhops</i> | 1 | 1 | VU |
| | | <i>Leptosittaca branickii</i> | 4 | 4 | VU |
| | | <i>Loriculus amabilis</i> | 2 | 2 | LC |
| | | <i>Loriculus beryllinus</i> | 3 | 3 | LC |
| | | <i>Loriculus catamene</i> | 0 | 0 | NT |
| | | <i>Loriculus exilis</i> | 0 | 0 | NT |
| | | <i>Loriculus flosculus</i> | 0 | 0 | EN |
| | | <i>Loriculus pusillus</i> | 4 | 4 | NT |
| | | <i>Loriculus stigmatus</i> | 4 | 4 | LC |
| | | <i>Loriculus vernalis</i> | 2 | 1 | LC |
| | | <i>Micropsitta geelvinkiana</i> | 2 | 2 | NT |
| | | <i>Micropsitta keiensis</i> | 4 | 4 | LC |
| | | <i>Micropsitta meeki</i> | 3 | 3 | LC |
| | | <i>Micropsitta pusio</i> | 4 | 4 | LC |
| | | <i>Nannopsittaca dachilleae</i> | 0 | 0 | NT |
| | | <i>Nannopsittaca panychlora</i> | 2 | 2 | LC |
| | | <i>Pionopsitta aurantiocephala</i> | 0 | 0 | |
| | | <i>Pionopsitta pyrrilia</i> | 0 | 0 | |

| Order | Family | Taxon | No. trade records | | IUCN |
|------------|---------|---------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Pionopsitta vulturina</i> | 1 | 1 | |
| | | <i>Platycercus venustus</i> | 2 | 2 | LC |
| | | <i>Poicephalus crassus</i> | 4 | 4 | LC |
| | | <i>Poicephalus senegalus versteri</i> | 0 | 0 | LC |
| | | <i>Prioniturus discurus</i> | 4 | 4 | LC |
| | | <i>Prioniturus flavicans</i> | 1 | 1 | NT |
| | | <i>Prioniturus luconensis</i> | 1 | 1 | VU |
| | | <i>Prioniturus mada</i> | 2 | 2 | LC |
| | | <i>Prioniturus montanus</i> | 2 | 2 | NT |
| | | <i>Prioniturus platenae</i> | 1 | 1 | VU |
| | | <i>Prioniturus verticalis</i> | 0 | 0 | EN |
| | | <i>Prosopieia personata</i> | 1 | 0 | NT |
| | | <i>Prosopieia splendens</i> | 0 | 0 | VU |
| | | <i>Prosopieia tabuensis</i> | 4 | 2 | LC |
| | | <i>Psittacella brehmii</i> | 4 | 3 | LC |
| | | <i>Psittacella madaraszi</i> | 0 | 0 | LC |
| | | <i>Psittacella modesta</i> | 0 | 0 | LC |
| | | <i>Psittacella picta</i> | 0 | 0 | LC |
| | | <i>Psittacula caniceps</i> | 0 | 0 | NT |
| | | <i>Psittacula finschii</i> | 2 | 2 | LC |
| | | <i>Psittacula intermedia</i> | 0 | 0 | |
| | | <i>Psittaculirostris edwardsii</i> | 5 | 1 | LC |
| | | <i>Psittaculirostris salvadorii</i> | 4 | 2 | VU |
| | | <i>Pyrrhura albipectus</i> | 2 | 2 | VU |
| | | <i>Pyrrhura caeruleiceps</i> | 0 | 0 | |
| | | <i>Pyrrhura calliptera</i> | 0 | 0 | VU |
| | | <i>Pyrrhura devillei</i> | 1 | 0 | NT |
| | | <i>Pyrrhura eisenmanni</i> | 0 | 0 | |
| | | <i>Pyrrhura hoematotis</i> | 1 | 0 | LC |
| | | <i>Pyrrhura lepida</i> | 4 | 1 | NT |
| | | <i>Pyrrhura orcesi</i> | 0 | 0 | EN |
| | | <i>Pyrrhura parvifrons</i> | 0 | 0 | |
| | | <i>Pyrrhura pfrimeri</i> | 0 | 0 | EN |
| | | <i>Pyrrhura picta picta</i> | 0 | 0 | |
| | | <i>Pyrrhura subandina</i> | 0 | 0 | |
| | | <i>Pyrrhura viridicata</i> | 0 | 0 | EN |
| | | <i>Tanygnathus gramineus</i> | 0 | 0 | VU |
| | | <i>Tanygnathus sumatranus</i> | 5 | 4 | LC |
| | | <i>Touit costaricensis</i> | 0 | 0 | VU |
| | | <i>Touit dilectissimus</i> | 3 | 3 | LC |
| | | <i>Touit melanonotus</i> | 2 | 2 | EN |
| | | <i>Touit purpuratus</i> | 3 | 3 | LC |
| | | <i>Touit stictopterus</i> | 2 | 2 | VU |
| | | <i>Touit surdus</i> | 0 | 0 | VU |
| | | <i>Tricharia malachitacea</i> | 2 | 1 | NT |
| RHEIFORMES | RHEIDAE | <i>Pterocnemis pennata pennata</i> | 3 | 0 | NT |

| Order | Family | Taxon | No. trade records | | IUCN |
|--------------|-----------|---------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| STRIGIFORMES | STRIGIDAE | <i>Aegolius ridgwayi</i> | 1 | 1 | LC |
| | | <i>Asio abyssinicus</i> | 3 | 3 | LC |
| | | <i>Asio madagascariensis</i> | 0 | 0 | LC |
| | | <i>Asio stygius</i> | 4 | 4 | LC |
| | | <i>Athene brama</i> | 0 | 0 | LC |
| | | <i>Bubo blakistoni</i> | 0 | 0 | |
| | | <i>Bubo cinerascens</i> | 4 | 4 | |
| | | <i>Bubo coromandus</i> | 0 | 0 | LC |
| | | <i>Bubo leucostictus</i> | 0 | 0 | LC |
| | | <i>Bubo nipalensis</i> | 3 | 2 | LC |
| | | <i>Bubo philippensis</i> | 2 | 2 | VU |
| | | <i>Bubo shelleyi</i> | 0 | 0 | NT |
| | | <i>Bubo sumatranus</i> | 1 | 0 | LC |
| | | <i>Bubo vosseleri</i> | 0 | 0 | VU |
| | | <i>Ciccaba huhula</i> | 5 | 3 | LC |
| | | <i>Ciccaba nigrolineata</i> | 0 | 0 | LC |
| | | <i>Glaucidium bolivianum</i> | 1 | 1 | LC |
| | | <i>Glaucidium capense</i> | 5 | 5 | LC |
| | | <i>Glaucidium castanonotum</i> | 0 | 0 | NT |
| | | <i>Glaucidium castanopterum</i> | 0 | 0 | LC |
| | | <i>Glaucidium costaricanum</i> | 0 | 0 | LC |
| | | <i>Glaucidium griseiceps</i> | 0 | 0 | LC |
| | | <i>Glaucidium hardyi</i> | 3 | 1 | LC |
| | | <i>Glaucidium minutissimum</i> | 1 | 1 | LC |
| | | <i>Glaucidium mooreorum</i> | 0 | 0 | CR |
| | | <i>Glaucidium nubicola</i> | 0 | 0 | VU |
| | | <i>Glaucidium palmarum</i> | 0 | 0 | LC |
| | | <i>Glaucidium parkeri</i> | 2 | 2 | LC |
| | | <i>Glaucidium peruanum</i> | 3 | 3 | LC |
| | | <i>Glaucidium radiatum</i> | 1 | 1 | LC |
| | | <i>Glaucidium sanchezi</i> | 0 | 0 | LC |
| | | <i>Glaucidium siju</i> | 0 | 0 | LC |
| | | <i>Glaucidium sjostedti</i> | 2 | 2 | LC |
| | | <i>Glaucidium tephronotum</i> | 3 | 3 | LC |
| | | <i>Gymnogaix lawrencii</i> | 0 | 0 | LC |
| | | <i>Jubula lettii</i> | 0 | 0 | DD |
| | | <i>Ketupa flavipes</i> | 0 | 0 | LC |
| | | <i>Ketupa zeylonensis</i> | 2 | 2 | LC |
| | | <i>Lophostrix cristata</i> | 1 | 1 | LC |
| | | <i>Micrathene whitneyi</i> | 2 | 2 | LC |
| | | <i>Nesasio solomonensis</i> | 3 | 3 | VU |
| | | <i>Ninox affinis</i> | 0 | 0 | NT |
| | | <i>Ninox boobook</i> | 5 | 5 | |
| | | <i>Ninox burhani</i> | 0 | 0 | NT |
| | | <i>Ninox connivens</i> | 0 | 0 | LC |
| | | <i>Ninox ios</i> | 0 | 0 | VU |
| | | <i>Ninox jacquinoti</i> | 3 | 3 | LC |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|--------|------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Ninox meeki</i> | 0 | 0 | LC |
| | | <i>Ninox ochracea</i> | 0 | 0 | NT |
| | | <i>Ninox odiosa</i> | 0 | 0 | VU |
| | | <i>Ninox punctulata</i> | 0 | 0 | LC |
| | | <i>Ninox rudolfi</i> | 0 | 0 | NT |
| | | <i>Ninox rufa</i> | 1 | 1 | LC |
| | | <i>Ninox squamipila</i> | 0 | 0 | LC |
| | | <i>Ninox strenua</i> | 4 | 4 | LC |
| | | <i>Ninox sumbaensis</i> | 0 | 0 | NT |
| | | <i>Ninox superciliaris</i> | 5 | 5 | LC |
| | | <i>Ninox theomacha</i> | 2 | 2 | LC |
| | | <i>Ninox variegata</i> | 0 | 0 | LC |
| | | <i>Otus albobularis</i> | 5 | 5 | LC |
| | | <i>Otus alfredi</i> | 0 | 0 | EN |
| | | <i>Otus alius</i> | 0 | 0 | DD |
| | | <i>Otus angelinae</i> | 0 | 0 | VU |
| | | <i>Otus atricapilla</i> | 0 | 0 | LC |
| | | <i>Otus balli</i> | 0 | 0 | NT |
| | | <i>Otus barbarus</i> | 3 | 3 | NT |
| | | <i>Otus beccarii</i> | 0 | 0 | EN |
| | | <i>Otus brookii</i> | 0 | 0 | LC |
| | | <i>Otus capnodes</i> | 0 | 0 | CR |
| | | <i>Otus clarkii</i> | 0 | 0 | LC |
| | | <i>Otus collari</i> | 0 | 0 | LC |
| | | <i>Otus colombianus</i> | 0 | 0 | NT |
| | | <i>Otus cooperi</i> | 2 | 2 | LC |
| | | <i>Otus elegans</i> | 3 | 3 | NT |
| | | <i>Otus fuliginosus</i> | 2 | 2 | NT |
| | | <i>Otus guatemalae</i> | 1 | 0 | LC |
| | | <i>Otus hartlaubi</i> | 0 | 0 | VU |
| | | <i>Otus hoyi</i> | 2 | 2 | |
| | | <i>Otus icterorhynchus</i> | 0 | 0 | LC |
| | | <i>Otus insularis</i> | 0 | 0 | EN |
| | | <i>Otus ireneae</i> | 3 | 3 | EN |
| | | <i>Otus koepckeae</i> | 1 | 1 | LC |
| | | <i>Otus madagascariensis</i> | 1 | 1 | |
| | | <i>Otus magicus</i> | 1 | 1 | LC |
| | | <i>Otus manadensis</i> | 1 | 1 | LC |
| | | <i>Otus mantananensis</i> | 1 | 1 | NT |
| | | <i>Otus marshalli</i> | 1 | 0 | NT |
| | | <i>Otus mayottensis</i> | 0 | 0 | LC |
| | | <i>Otus mentawi</i> | 0 | 0 | NT |
| | | <i>Otus mindorensis</i> | 0 | 0 | NT |
| | | <i>Otus mirus</i> | 2 | 2 | NT |
| | | <i>Otus moheliensis</i> | 0 | 0 | CR |
| | | <i>Otus nudipes</i> | 0 | 0 | LC |
| | | <i>Otus pauliani</i> | 0 | 0 | CR |

| Order | Family | Taxon | No. trade records | | IUCN |
|-----------------------|-----------|----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Otus pambaensis</i> | 0 | 0 | VU |
| | | <i>Otus petersoni</i> | 0 | 0 | LC |
| | | <i>Otus rufescens</i> | 3 | 2 | NT |
| | | <i>Otus sagittatus</i> | 1 | 1 | VU |
| | | <i>Otus sanctaecatrinae</i> | 0 | 0 | LC |
| | | <i>Otus seductus</i> | 0 | 0 | NT |
| | | <i>Otus senegalensis</i> | 4 | 2 | LC |
| | | <i>Otus silvicola</i> | 0 | 0 | LC |
| | | <i>Otus spilocephalus</i> | 2 | 1 | LC |
| | | <i>Otus thilohoffmanni</i> | 0 | 0 | EN |
| | | <i>Otus umbra</i> | 0 | 0 | NT |
| | | <i>Otus vermiculatus</i> | 4 | 4 | |
| | | <i>Pseudoscops grammicus</i> | 2 | 2 | LC |
| | | <i>Ptilopsis granti</i> | 1 | 1 | |
| | | <i>Pulsatrix koenigswaldiana</i> | 1 | 1 | LC |
| | | <i>Pyroglaux podargina</i> | 0 | 0 | |
| | | <i>Sceloglaux albifacies</i> | 0 | 0 | EX |
| | | <i>Scotopelia ussheri</i> | 0 | 0 | EN |
| | | <i>Strix albitarsis</i> | 1 | 1 | LC |
| | | <i>Strix butleri</i> | 1 | 1 | LC |
| | | <i>Strix chacoensis</i> | 0 | 0 | LC |
| | | <i>Strix fulvescens</i> | 2 | 2 | LC |
| | | <i>Strix hylophila</i> | 0 | 0 | NT |
| | | <i>Strix leptogrammica</i> | 1 | 0 | LC |
| | | <i>Strix ocellata</i> | 0 | 0 | LC |
| | | <i>Strix uralensis davidi</i> | 0 | 0 | |
| | | <i>Uroglaux dimorpha</i> | 1 | 1 | DD |
| | | <i>Xenoglaux loweryi</i> | 1 | 0 | EN |
| | TYTONIDAE | <i>Phodilus prigoginei</i> | 0 | 0 | EN |
| | | <i>Tyto aurantia</i> | 0 | 0 | VU |
| | | <i>Tyto capensis</i> | 5 | 5 | LC |
| | | <i>Tyto glaucops</i> | 4 | 4 | LC |
| | | <i>Tyto inexpectata</i> | 0 | 0 | VU |
| | | <i>Tyto manusi</i> | 0 | 0 | VU |
| | | <i>Tyto multipunctata</i> | 0 | 0 | |
| | | <i>Tyto nigrobrunnea</i> | 0 | 0 | EN |
| | | <i>Tyto novaehollandiae</i> | 2 | 2 | LC |
| | | <i>Tyto rosenbergii</i> | 0 | 0 | LC |
| | | <i>Tyto sororcula</i> | 0 | 0 | DD |
| | | <i>Tyto tenebricosa</i> | 2 | 2 | LC |
| CLASS REPTILIA | | | | | |
| SAURIA | AGAMIDAE | <i>Uromastix alfredschmidti</i> | 0 | 0 | NT |
| | | <i>Uromastix asmussi</i> | 3 | 2 | |
| | | <i>Uromastix leptieni</i> | 1 | 0 | |
| | | <i>Uromastix loricata</i> | 0 | 0 | LC |
| | | <i>Uromastix macfadyeni</i> | 0 | 0 | |
| | | <i>Uromastix occidentalis</i> | 0 | 0 | |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|----------------|-----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Uromastyx thomasi</i> | 2 | 1 | |
| | | <i>Uromastyx yemenensis</i> | 0 | 0 | |
| | CHAMAELEONIDAE | <i>Bradypodion atromontanum</i> | 0 | 0 | |
| | | <i>Bradypodion caeruleogula</i> | 0 | 0 | |
| | | <i>Bradypodion cafferum</i> | 1 | 1 | EN |
| | | <i>Bradypodion damaranum</i> | 2 | 2 | |
| | | <i>Bradypodion dracomontanum</i> | 5 | 5 | LC |
| | | <i>Bradypodion gutturale</i> | 0 | 0 | |
| | | <i>Bradypodion nkandlae</i> | 0 | 0 | |
| | | <i>Bradypodion pumilum</i> | 5 | 4 | EN |
| | | <i>Bradypodion setaroi</i> | 4 | 4 | LC |
| | | <i>Bradypodion transvaalense</i> | 5 | 5 | LC |
| | | <i>Bradypodion ventrale</i> | 1 | 1 | LC |
| | | <i>Calumma ambreense</i> | 0 | 0 | |
| | | <i>Calumma linota</i> | 0 | 0 | |
| | | <i>Calumma marojezense</i> | 0 | 0 | |
| | | <i>Chamaeleo affinis</i> | 3 | 3 | |
| | | <i>Chamaeleo arabicus</i> | 0 | 0 | |
| | | <i>Chamaeleo balebicornutus</i> | 0 | 0 | |
| | | <i>Chamaeleo camerunensis</i> | 4 | 4 | |
| | | <i>Chamaeleo chapini</i> | 0 | 0 | |
| | | <i>Chamaeleo conirostratus</i> | 0 | 0 | |
| | | <i>Chamaeleo harenae</i> | 0 | 0 | |
| | | <i>Chamaeleo ituriensis</i> | 2 | 2 | |
| | | <i>Chamaeleo kinetensis</i> | 1 | 1 | |
| | | <i>Chamaeleo laevigatus</i> | 3 | 2 | |
| | | <i>Chamaeleo marsabitensis</i> | 0 | 0 | |
| | | <i>Chamaeleo monachus</i> | 2 | 0 | |
| | | <i>Chamaeleo namaquensis</i> | 2 | 2 | |
| | | <i>Chamaeleo narraioaca</i> | 0 | 0 | |
| | | <i>Chamaeleo necasi</i> | 0 | 0 | |
| | | <i>Chamaeleo ntunte</i> | 0 | 0 | |
| | | <i>Chamaeleo schoutedeni</i> | 2 | 2 | |
| | | <i>Chamaeleo schubotzi</i> | 0 | 0 | |
| | | <i>Chamaeleo tremperi</i> | 0 | 0 | |
| | | <i>Chamaeleo zeylanicus</i> | 5 | 5 | |
| | | <i>Furcifer tuzetae</i> | 1 | 1 | |
| | | <i>Kinyongia boehmei</i> | 0 | 0 | |
| | | <i>Kinyongia excubitor</i> | 0 | 0 | |
| | | <i>Kinyongia matschiei</i> | 0 | 0 | |
| | | <i>Kinyongia multituberculata</i> | 0 | 0 | |
| | | <i>Kinyongia uluguruensis</i> | 0 | 0 | |
| | | <i>Kinyongia vosseleri</i> | 0 | 0 | |
| | | <i>Nadzikambia mlanjense</i> | 0 | 0 | |
| | CORDYLIDAE | <i>Cordylus angolensis</i> | 0 | 0 | |
| | | <i>Cordylus aridus</i> | 0 | 0 | EN |
| | | <i>Cordylus barbertonensis</i> | 2 | 2 | |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|------------|------------------------------------|--------------------------|------|-------|
| | | | All sources except C & D | Wild | |
| | | <i>Cordylus beraduccii</i> | 4 | 3 | |
| | | <i>Cordylus breyeri</i> | 2 | 2 | |
| | | <i>Cordylus campbelli</i> | 2 | 2 | DD |
| | | <i>Cordylus capensis</i> | 1 | 1 | |
| | | <i>Cordylus cataphractus</i> | 4 | 1 | VU |
| | | <i>Cordylus cloetei</i> | 0 | 0 | |
| | | <i>Cordylus coeruleopunctatus</i> | 1 | 1 | |
| | | <i>Cordylus cordylus</i> | 3 | 2 | |
| | | <i>Cordylus depressus</i> | 3 | 2 | |
| | | <i>Cordylus fasciatus</i> | 0 | 0 | |
| | | <i>Cordylus imkae</i> | 3 | 3 | |
| | | <i>Cordylus jonesii</i> | 0 | 0 | |
| | | <i>Cordylus jordani</i> | 2 | 2 | |
| | | <i>Cordylus langi</i> | 1 | 1 | LR/nt |
| | | <i>Cordylus lawrenci</i> | 0 | 0 | LR/nt |
| | | <i>Cordylus machadoi</i> | 0 | 0 | |
| | | <i>Cordylus macropholis</i> | 2 | 2 | |
| | | <i>Cordylus mclachlani</i> | 0 | 0 | VU |
| | | <i>Cordylus meculae</i> | 0 | 0 | EN |
| | | <i>Cordylus melanotus</i> | 5 | 2 | |
| | | <i>Cordylus microlepidotus</i> | 0 | 0 | |
| | | <i>Cordylus minor</i> | 0 | 0 | |
| | | <i>Cordylus namaquensis</i> | 1 | 1 | |
| | | <i>Cordylus nebulosus</i> | 0 | 0 | VU |
| | | <i>Cordylus niger</i> | 3 | 3 | |
| | | <i>Cordylus nyikae</i> | 0 | 0 | |
| | | <i>Cordylus oelofseni</i> | 1 | 1 | |
| | | <i>Cordylus peersi</i> | 1 | 1 | |
| | | <i>Cordylus polyzonus</i> | 1 | 1 | |
| | | <i>Cordylus pustulatus</i> | 1 | 1 | |
| | | <i>Cordylus regius</i> | 0 | 0 | |
| | | <i>Cordylus rivaie</i> | 0 | 0 | LC |
| | | <i>Cordylus spinosus</i> | 1 | 1 | LC |
| | | <i>Cordylus subviridis</i> | 0 | 0 | |
| | | <i>Cordylus tasmani</i> | 1 | 1 | VU |
| | | <i>Cordylus transvaalensis</i> | 3 | 2 | |
| | | <i>Cordylus vandami</i> | 4 | 3 | |
| | GEKKONIDAE | <i>Cyrtodactylus serpensinsula</i> | 1 | 1 | |
| | | <i>Phelsuma andamanense</i> | 0 | 0 | LC |
| | | <i>Phelsuma astriata</i> | 0 | 0 | LC |
| | | <i>Phelsuma borbonica</i> | 1 | 1 | |
| | | <i>Phelsuma edwardnewtonii</i> | 0 | 0 | |
| | | <i>Phelsuma gigas</i> | 0 | 0 | EX |
| | | <i>Phelsuma grandis</i> | 0 | 0 | |
| | | <i>Phelsuma inexpectata</i> | 1 | 1 | |
| | | <i>Phelsuma kochi</i> | 0 | 0 | |
| | | <i>Phelsuma ocellata</i> | 0 | 0 | LR/nt |

| Order | Family | Taxon | No. trade records | | IUCN |
|-------|------------|----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Phelsuma pasteuri</i> | 3 | 3 | |
| | | <i>Phelsuma rosagularis</i> | 2 | 2 | |
| | | <i>Phelsuma sundbergi</i> | 1 | 0 | LC |
| | IGUANIDAE | <i>Phrynosoma blainvillii</i> | 3 | 1 | |
| | | <i>Phrynosoma cerroense</i> | 0 | 0 | |
| | | <i>Phrynosoma wigginsii</i> | 0 | 0 | |
| | LACERTIDAE | <i>Podarcis lilfordi</i> | 0 | 0 | EN |
| | | <i>Podarcis pityusensis</i> | 1 | 0 | NT |
| | TEIIDAE | <i>Crocodylurus amazonicus</i> | 3 | 3 | LC |
| | | <i>Dracaena paraguayensis</i> | 0 | 0 | |
| | | <i>Tupinambis cerradensis</i> | 1 | 1 | |
| | | <i>Tupinambis duseni</i> | 0 | 0 | |
| | | <i>Tupinambis longilineus</i> | 1 | 1 | |
| | | <i>Tupinambis palustris</i> | 0 | 0 | |
| | | <i>Tupinambis quadrilineatus</i> | 2 | 0 | |
| | VARANIDAE | <i>Varanus auffenbergi</i> | 0 | 0 | |
| | | <i>Varanus baritji</i> | 2 | 2 | |
| | | <i>Varanus bogerti</i> | 2 | 2 | |
| | | <i>Varanus brevicauda</i> | 1 | 1 | |
| | | <i>Varanus bushi</i> | 0 | 0 | |
| | | <i>Varanus caerulivirens</i> | 0 | 0 | |
| | | <i>Varanus caudolineatus</i> | 0 | 0 | |
| | | <i>Varanus cerambonensis</i> | 0 | 0 | |
| | | <i>Varanus cumingi</i> | 0 | 0 | LC |
| | | <i>Varanus eremius</i> | 1 | 1 | |
| | | <i>Varanus finschi</i> | 5 | 4 | LC |
| | | <i>Varanus giganteus</i> | 4 | 3 | |
| | | <i>Varanus gilleni</i> | 1 | 0 | |
| | | <i>Varanus glauerti</i> | 3 | 1 | LC |
| | | <i>Varanus glebopalma</i> | 2 | 2 | |
| | | <i>Varanus juxtindicus</i> | 0 | 0 | |
| | | <i>Varanus keithhornei</i> | 0 | 0 | |
| | | <i>Varanus kingorum</i> | 4 | 2 | |
| | | <i>Varanus kordensis</i> | 1 | 0 | |
| | | <i>Varanus mabitang</i> | 1 | 1 | EN |
| | | <i>Varanus marmoratus</i> | 1 | 1 | LC |
| | | <i>Varanus mertensi</i> | 2 | 0 | |
| | | <i>Varanus mitchelli</i> | 0 | 0 | |
| | | <i>Varanus nuchalis</i> | 0 | 0 | NT |
| | | <i>Varanus olivaceus</i> | 5 | 1 | VU |
| | | <i>Varanus pilbarensis</i> | 4 | 1 | |
| | | <i>Varanus primordius</i> | 2 | 2 | LC |
| | | <i>Varanus rainerguentheri</i> | 0 | 0 | |
| | | <i>Varanus rosenbergi</i> | 3 | 2 | LC |
| | | <i>Varanus scalaris</i> | 0 | 0 | LC |
| | | <i>Varanus semiremex</i> | 2 | 2 | |
| | | <i>Varanus spenceri</i> | 1 | 1 | |

| Order | Family | Taxon | No. trade records | | IUCN |
|-----------|----------------|----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Varanus telenesetes</i> | 1 | 1 | DD |
| | | <i>Varanus togianus</i> | 0 | 0 | |
| | | <i>Varanus tristis</i> | 5 | 1 | |
| | | <i>Varanus varius</i> | 4 | 3 | |
| | | <i>Varanus yemenensis</i> | 1 | 1 | DD |
| | | <i>Varanus zugorum</i> | 0 | 0 | |
| SERPENTES | BOIDAE | <i>Corallus cookii</i> | 5 | 5 | |
| | | <i>Corallus cropanii</i> | 0 | 0 | EN |
| | | <i>Corallus grenadensis</i> | 0 | 0 | |
| | | <i>Corallus ruschenbergerii</i> | 5 | 2 | |
| | | <i>Epicrates chrysogaster</i> | 5 | 4 | |
| | | <i>Epicrates exsul</i> | 1 | 1 | |
| | | <i>Epicrates fordii</i> | 4 | 4 | |
| | | <i>Epicrates gracilis</i> | 2 | 2 | |
| | | <i>Eryx elegans</i> | 0 | 0 | |
| | | <i>Eryx jaculus</i> | 3 | 2 | |
| | | <i>Eryx johnii</i> | 4 | 3 | |
| | | <i>Eryx somalicus</i> | 0 | 0 | |
| | | <i>Eryx whitakeri</i> | 0 | 0 | |
| | | <i>Eunectes beniensis</i> | 0 | 0 | LC |
| | | <i>Eunectes deschauenseei</i> | 0 | 0 | DD |
| | COLUBRIDAE | <i>Elachistodon westermanni</i> | 2 | 2 | LC |
| | ELAPIDAE | <i>Hoplocephalus bungaroides</i> | 0 | 0 | VU |
| | | <i>Naja mandalayensis</i> | 0 | 0 | |
| | | <i>Naja philippinensis</i> | 5 | 2 | NT |
| | | <i>Naja sagittifera</i> | 0 | 0 | |
| | | <i>Naja samarensis</i> | 3 | 3 | LC |
| | PYTHONIDAE | <i>Antaresia maculosa</i> | 5 | 0 | |
| | | <i>Antaresia perthensis</i> | 0 | 0 | |
| | | <i>Antaresia stimsoni</i> | 1 | 0 | |
| | | <i>Morelia carinata</i> | 1 | 1 | |
| | | <i>Morelia clastolepis</i> | 0 | 0 | |
| | | <i>Morelia kinghorni</i> | 0 | 0 | |
| | | <i>Morelia nauta</i> | 0 | 0 | |
| | | <i>Morelia oenpelliensis</i> | 0 | 0 | |
| | | <i>Morelia tracyae</i> | 0 | 0 | |
| | | <i>Python natalensis</i> | 0 | 0 | |
| | TROPIDOPHIIDAE | <i>Exiliboa placata</i> | 2 | 2 | VU |
| | | <i>Trachyboa boulengeri</i> | 2 | 1 | |
| | | <i>Trachyboa gularis</i> | 0 | 0 | |
| | | <i>Tropidophis battersbyi</i> | 0 | 0 | |
| | | <i>Tropidophis canus</i> | 1 | 1 | |
| | | <i>Tropidophis caymanensis</i> | 0 | 0 | |
| | | <i>Tropidophis celiae</i> | 0 | 0 | |
| | | <i>Tropidophis feicki</i> | 0 | 0 | |
| | | <i>Tropidophis fuscus</i> | 0 | 0 | |
| | | <i>Tropidophis greenwayi</i> | 4 | 4 | |

| Order | Family | Taxon | No. trade records | | IUCN |
|-----------------------------|----------------|-----------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| | | <i>Tropidophis hendersoni</i> | 0 | 0 | CR |
| | | <i>Tropidophis maculatus</i> | 0 | 0 | |
| | | <i>Tropidophis morenoi</i> | 0 | 0 | |
| | | <i>Tropidophis nigriventris</i> | 0 | 0 | |
| | | <i>Tropidophis pardalis</i> | 0 | 0 | LC |
| | | <i>Tropidophis paucisquamis</i> | 0 | 0 | |
| | | <i>Tropidophis pilsbryi</i> | 0 | 0 | |
| | | <i>Tropidophis semicinctus</i> | 0 | 0 | |
| | | <i>Tropidophis spiritus</i> | 0 | 0 | |
| | | <i>Tropidophis taczanowskyi</i> | 2 | 1 | |
| | | <i>Tropidophis wrighti</i> | 0 | 0 | |
| | | <i>Tropidophis xanthogaster</i> | 0 | 0 | |
| | | <i>Ungaliophis continentalis</i> | 0 | 0 | |
| | | <i>Ungaliophis panamensis</i> | 1 | 1 | |
| | VIPERIDAE | <i>Vipera wagneri</i> | 1 | 1 | CR |
| TESTUDINES | PODOCNEMIDIDAE | <i>Peltocephalus dumerilianus</i> | 0 | 0 | VU |
| | | <i>Podocnemis lewyana</i> | 2 | 2 | EN |
| | | <i>Podocnemis vogli</i> | 4 | 2 | |
| | TRIONYCHIDAE | <i>Lissemys scutata</i> | 0 | 0 | DD |
| CLASS ACTINOPTERYGII | | | | | |
| CYPRINIFORMES | CYPRINIDAE | <i>Caecobarbus geertsii</i> | 1 | 1 | VU |
| PHYLUM ARTHROPODA | | | | | |
| CLASS ARACHNIDA | | | | | |
| ARANEAE | THERAPHOSIDAE | <i>Aphonopelma albiceps</i> | 0 | 0 | |
| | | <i>Aphonopelma pallidum</i> | 0 | 0 | |
| | | <i>Brachypelma andrewi</i> | 2 | 2 | |
| | | <i>Brachypelma angustum</i> | 0 | 0 | |
| | | <i>Brachypelma annitha</i> | 0 | 0 | |
| | | <i>Brachypelma auratum</i> | 5 | 1 | |
| | | <i>Brachypelma aureoiceps</i> | 1 | 1 | |
| | | <i>Brachypelma baumgarteni</i> | 0 | 0 | |
| | | <i>Brachypelma embrithes</i> | 0 | 0 | |
| | | <i>Brachypelma epicureanum</i> | 1 | 1 | |
| | | <i>Brachypelma fossorium</i> | 3 | 3 | |
| | | <i>Brachypelma hamorii</i> | 0 | 0 | |
| | | <i>Brachypelma kahlenbergi</i> | 0 | 0 | |
| | | <i>Brachypelma klaasi</i> | 5 | 3 | |
| | | <i>Brachypelma ruhnaui</i> | 0 | 0 | |
| | | <i>Brachypelma sabulosum</i> | 1 | 1 | |
| | | <i>Brachypelma schroederi</i> | 0 | 0 | |
| | | <i>Brachypelma verdezi</i> | 0 | 0 | |
| SCORPIONES | SCORPIONIDAE | <i>Pandinus gambiensis</i> | 2 | 1 | |

| Order | Family | Taxon | No. trade records | | IUCN |
|---------------------------|--------------|-------------------------------------|--------------------------|------|------|
| | | | All sources except C & D | Wild | |
| CLASS INSECTA | | | | | |
| LEPIDOPTERA | PAPILIONIDAE | <i>Bhutanitis ludlowi</i> | 0 | 0 | VU |
| | | <i>Ornithoptera akakeae</i> | 0 | 0 | |
| | | <i>Parnassius apollo apollo</i> | 0 | 0 | VU |
| | | <i>Teinopalpus aureus</i> | 2 | 0 | DD |
| | | <i>Troides darsius</i> | 4 | 0 | |
| | | <i>Troides minos</i> | 2 | 2 | |
| PHYLUM ANNELIDA | | | | | |
| CLASS HIRUDINOIDEA | | | | | |
| ARHYNCHOBDELLIDA | HIRUDINIDAE | <i>Hirudo verbana</i> | 0 | 0 | |
| PHYLUM MOLLUSCA | | | | | |
| CLASS BIVALVIA | | | | | |
| UNIONOIDA | UNIONIDAE | <i>Epioblasma torulosa rangiana</i> | 1 | 1 | |
| VENEROIDA | TRIDACNIDAE | <i>Tridacna rosewateri</i> | 0 | 0 | VU |
| | | <i>Tridacna tevoroa</i> | 1 | 1 | VU |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |